

Product datasheet for RC209120L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PLC delta 3 (PLCD3) (NM_133373) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PLC delta 3 (PLCD3) (NM_133373) Human Tagged ORF Clone Lentiviral Particle

Symbol: PLC delta 3
Synonyms: PLC-delta-3
Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_133373 **ORF Size:** 2367 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209120).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 133373.3, NP 588614.1

 RefSeq Size:
 3477 bp

 RefSeq ORF:
 2370 bp

 Locus ID:
 113026

 UniProt ID:
 Q8N3E9

 Cytogenetics:
 17q21.31

Domains: C2, PI-PLC-X, PI-PLC-Y, PH

Protein Families: Druggable Genome





PLC delta 3 (PLCD3) (NM_133373) Human Tagged ORF Clone Lentiviral Particle - RC209120L4V

Protein Pathways: Calcium signaling pathway, Inositol phosphate metabolism, Metabolic pathways,

Phosphatidylinositol signaling system

MW: 89.1 kDa

Gene Summary: This gene encodes a member of the phospholipase C family, which catalyze the hydrolysis of

phosphatidylinositol 4,5-bisphosphate to generate the second messengers diacylglycerol and

inositol 1,4,5-trisphosphate (IP3). Diacylglycerol and IP3 mediate a variety of cellular

responses to extracellular stimuli by inducing protein kinase C and increasing cytosolic Ca(2+) concentrations. This enzyme localizes to the plasma membrane and requires calcium for activation. Its activity is inhibited by spermine, sphingosine, and several phospholipids.

[provided by RefSeq, Jul 2008]