

Product datasheet for RC209128L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DUSP5 (NM_004419) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: DUSP5 (NM_004419) Human Tagged ORF Clone Lentiviral Particle

Symbol: DUSP5

Synonyms: DUSP; HVH3

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_004419

ORF Size: 1152 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC209128).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 004419.3</u>

 RefSeq Size:
 2545 bp

 RefSeq ORF:
 1155 bp

 Locus ID:
 1847

 UniProt ID:
 Q16690

 Cytogenetics:
 10q25.2

Domains: DSPc, RHOD, PTPc_motif

Protein Families: Phosphatase





Protein Pathways: MAPK signaling pathway

MW: 42.1 kDa

Gene Summary: The protein encoded by this gene is a member of the dual specificity protein phosphatase

subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK1, is expressed in a variety of tissues with the highest levels in pancreas and brain, and is localized in the nucleus.

[provided by RefSeq, Jul 2008]