

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC209148L4V

## SCD1 (SCD) (NM\_005063) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	SCD1 (SCD) (NM_005063) Human Tagged ORF Clone Lentiviral Particle
Symbol:	SCD1
Synonyms:	FADS5; hSCD1; MSTP008; SCD1; SCDOS
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_005063
ORF Size:	1077 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC209148).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 005063.4</u>
RefSeq Size:	5473 bp
RefSeq ORF:	1080 bp
Locus ID:	6319
UniProt ID:	<u>000767</u>
Cytogenetics:	10q24.31
Domains:	FA_desaturase
Protein Families:	Transmembrane



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	SCD1 (SCD) (NM_005063) Human Tagged ORF Clone Lentiviral Particle – RC209148L4V
Protein Pathways	Biosynthesis of unsaturated fatty acids, PPAR signaling pathway
MW:	41.5 kDa
Gene Summary:	This gene encodes an enzyme involved in fatty acid biosynthesis, primarily the synthesis of oleic acid. The protein belongs to the fatty acid desaturase family and is an integral membrane protein located in the endoplasmic reticulum. Transcripts of approximately 3.9 and 5.2 kb, differing only by alternative polyadenlyation signals, have been detected. A gene encoding a similar enzyme is located on chromosome 4 and a pseudogene of this gene is located on chromosome 17. [provided by RefSeq, Sep 2015]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US