

Product datasheet for RC209150L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IGFBP3 (NM_000598) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IGFBP3 (NM_000598) Human Tagged ORF Clone Lentiviral Particle

Symbol: IGFBP3

Synonyms: BP-53; IBP3

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM 000598

ORF Size: 873 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209150).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000598.4

 RefSeq Size:
 2620 bp

 RefSeq ORF:
 876 bp

 Locus ID:
 3486

 UniProt ID:
 P17936

 Cytogenetics:
 7p12.3

Domains: thyroglobulin_1, IB

Protein Families: Druggable Genome, Secreted Protein





IGFBP3 (NM_000598) Human Tagged ORF Clone Lentiviral Particle - RC209150L4V

Protein Pathways: p53 signaling pathway

MW: 31.67 kDa

Gene Summary: This gene is a member of the insulin-like growth factor binding protein (IGFBP) family and

encodes a protein with an IGFBP domain and a thyroglobulin type-I domain. The protein forms a ternary complex with insulin-like growth factor acid-labile subunit (IGFALS) and either insulin-like growth factor (IGF) I or II. In this form, it circulates in the plasma, prolonging the

half-life of IGFs and altering their interaction with cell surface receptors. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

[provided by RefSeq, Jul 2008]