

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC209266L1V

Prolactin Receptor (PRLR) (NM_000949) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Prolactin Receptor (PRLR) (NM_000949) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Prolactin Receptor
Synonyms:	HPRL; hPRLrl; MFAB; RI-PRLR
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_000949
ORF Size:	1866 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC209266).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 000949.2</u>
RefSeq Size:	11694 bp
RefSeq ORF:	1869 bp
Locus ID:	5618
UniProt ID:	<u>P16471</u>
Cytogenetics:	5p13.2
Domains:	FN3
Protein Families:	Druggable Genome, Transmembrane



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Prolact RC2092	in Receptor (PRLR) (NM_000949) Human Tagged ORF Clone Lentiviral Particle – 66L1V
Protein Pathway	/s:	Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway, Neuroactive ligand- receptor interaction
MW:		69.5 kDa
Gene Summary:		This gene encodes a receptor for the anterior pituitary hormone, prolactin, and belongs to the type I cytokine receptor family. Prolactin-dependent signaling occurs as the result of ligand-induced dimerization of the prolactin receptor. Several alternatively spliced transcript variants encoding different membrane-bound and soluble isoforms have been described for this gene, which may function to modulate the endocrine and autocrine effects of prolactin in normal tissue and cancer. [provided by RefSeq, Feb 2011]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US