

## Product datasheet for RC209267L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Natriuretic Peptide Receptor A (NPR1) (NM\_000906) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Natriuretic Peptide Receptor A (NPR1) (NM\_000906) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Natriuretic Peptide Receptor A

Synonyms: ANPa; ANPRA; GUC2A; GUCY2A; NPRA

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_000906

ORF Size: 3183 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as (RC209267).

**OTI Disclaimer:** 

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 000906.2</u>, <u>NP 000897.2</u>

 RefSeq Size:
 4201 bp

 RefSeq ORF:
 3186 bp

 Locus ID:
 4881

 UniProt ID:
 P16066

Cytogenetics: 1q21.3

**Domains:** pkinase, CYCc, ANF\_receptor





**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Purine metabolism, Vascular smooth muscle contraction


**MW:** 119 kDa

**Gene Summary:** Guanylyl cyclases, catalyzing the production of cGMP from GTP, are classified as soluble and

membrane forms (Garbers and Lowe, 1994 [PubMed 7982997]). The membrane guanylyl cyclases, often termed guanylyl cyclases A through F, form a family of cell-surface receptors with a similar topographic structure: an extracellular ligand-binding domain, a single membrane-spanning domain, and an intracellular region that contains a protein kinase-like domain and a cyclase catalytic domain. GC-A and GC-B function as receptors for natriuretic peptides; they are also referred to as atrial natriuretic peptide receptor A (NPR1) and type B (NPR2; MIM 108961). Also see NPR3 (MIM 108962), which encodes a protein with only the ligand-binding transmembrane and 37-amino acid cytoplasmic domains. NPR1 is a membrane-bound guanylate cyclase that serves as the receptor for both atrial and brain natriuretic peptides (ANP (MIM 108780) and BNP (MIM 600295), respectively).[supplied by

OMIM, May 2009]

## **Product images:**



[RC209267L4] was used to prepare Lentiviral particles using [TR30037] packaging kit. HEK293T cells were transduced with RC209267L4V particle to overexpress human NPR1-mGFP fusion protein.