

Product datasheet for RC209272L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Factor D (CFD) (NM_001928) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Factor D (CFD) (NM_001928) Human Tagged ORF Clone Lentiviral Particle

Symbol: Factor D

Synonyms: ADIPSIN; ADN; DF; PFD

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 001928

Tag: Myc-DDK

ORF Size: 759 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209272).

Sequence:

ACCN:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001928.2

 RefSeq Size:
 1173 bp

 RefSeq ORF:
 762 bp

 Locus ID:
 1675

 UniProt ID:
 P00746

 Cytogenetics:
 19p13.3

Domains: Tryp_SPc

Protein Families: Druggable Genome, Protease, Secreted Protein







Protein Pathways: Complement and coagulation cascades

MW: 27 kDa

Gene Summary: This gene encodes a member of the S1, or chymotrypsin, family of serine peptidases. This

protease catalyzes the cleavage of factor B, the rate-limiting step of the alternative pathway of complement activation. This protein also functions as an adipokine, a cell signaling protein secreted by adipocytes, which regulates insulin secretion in mice. Mutations in this gene underlie complement factor D deficiency, which is associated with recurrent bacterial meningitis infections in human patients. Alternative splicing of this gene results in multiple

transcript variants. At least one of these variants encodes a preproprotein that is

proteolytically processed to generate the mature protease. [provided by RefSeq, Nov 2015]