

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC209297L3V

TEX19 (NM_207459) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	TEX19 (NM_207459) Human Tagged ORF Clone Lentiviral Particle
Symbol:	TEX19
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_207459
ORF Size:	492 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC209297).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 207459.1</u>
RefSeq Size:	1907 bp
RefSeq ORF:	495 bp
Locus ID:	400629
UniProt ID:	<u>Q8NA77</u>
Cytogenetics:	17q25.3
MW:	18.3 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Gene Summary:Required during spermatogenesis and placenta development, participating in the repression
of retrotransposable elements and prevent their mobilization. Collaborates with the Piwi-
interacting RNA (piRNA) pathway, which mediates the repression of transposable elements
during meiosis by forming complexes composed of piRNAs and Piwi proteins. Interacts with
Piwi proteins and directly binds piRNAs, a class of 24 to 30 nucleotide RNAs that are
generated by a Dicer-independent mechanism and are primarily derived from transposons
and other repeated sequence elements. Also during spermatogenesis, promotes, with UBR2,
SPO11-dependent recombination foci to accumulate and drive robust homologous
chromosome synapsis (By similarity). Interacts with LINE-1 retrotransposon encoded LIRE1,
stimulates LIRE1 polyubiquitination, mediated by UBR2, and degradation, inhibiting LINE-1
retranstoposon mobilization (PubMed:28806172).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US