

Product datasheet for RC209305L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SOCS3 (NM_003955) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SOCS3 (NM_003955) Human Tagged ORF Clone Lentiviral Particle

Symbol: SOCS3

Synonyms: ATOD4; CIS3; Cish3; SOCS-3; SSI-3; SSI3

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_003955

ORF Size: 675 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209305).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 003955.3

 RefSeq Size:
 2737 bp

 RefSeq ORF:
 678 bp

 Locus ID:
 9021

 UniProt ID:
 014543

 Cytogenetics:
 17q25.3

Domains: SH2, SOCS

Protein Families: Druggable Genome





SOCS3 (NM_003955) Human Tagged ORF Clone Lentiviral Particle - RC209305L4V

Protein Pathways: Adipocytokine signaling pathway, Insulin signaling pathway, Jak-STAT signaling pathway, Type

II diabetes mellitus, Ubiquitin mediated proteolysis

MW: 24.8 kDa

Gene Summary: This gene encodes a member of the STAT-induced STAT inhibitor (SSI), also known as

suppressor of cytokine signaling (SOCS), family. SSI family members are cytokine-inducible negative regulators of cytokine signaling. The expression of this gene is induced by various cytokines, including IL6, IL10, and interferon (IFN)-gamma. The protein encoded by this gene

can bind to JAK2 kinase, and inhibit the activity of JAK2 kinase. Studies of the mouse counterpart of this gene suggested the roles of this gene in the negative regulation of fetal

liver hematopoiesis, and placental development. [provided by RefSeq, Jul 2008]