

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC209308L3V

## RNF111 (NM\_017610) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Due du et Ture et	Lontinizal Devideo
Product Type:	Lentiviral Particles
Product Name:	RNF111 (NM_017610) Human Tagged ORF Clone Lentiviral Particle
Symbol:	RNF111
Synonyms:	ARK; hRNF111
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_017610
ORF Size:	2955 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC209308).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 017610.6</u>
RefSeq Size:	5902 bp
RefSeq ORF:	2961 bp
Locus ID:	54778
UniProt ID:	Q6ZNA4
Cytogenetics:	15q22.1-q22.2
Domains:	RING
Protein Families:	Druggable Genome



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	RNF111 (NM_017610) Human Tagged ORF Clone Lentiviral Particle – RC209308L3V
MW:	107.8 kDa
Gene Summary:	The protein encoded by this gene is a nuclear RING-domain containing E3 ubiquitin ligase. This protein interacts with the transforming growth factor (TGF) -beta/NODAL signaling pathway by promoting the ubiquitination and proteosomal degradation of negative regulators, like SMAD proteins, and thereby enhances TGF-beta target-gene transcription. As a modulator of the nodal signaling cascade, this gene plays a critical role in the induction of mesoderm during embryonic development. Alternative splicing of this gene results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2012]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US