

Product datasheet for **RC209326**

PSMB5 (NM_002797) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PSMB5 (NM_002797) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	PSMB5
Synonyms:	LMPX; MB1; X
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC209326 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCGTTGCCAGCGTGTGGAGAGACCGCTACCGGTGAACCAGCGGGTTTTTCGGACTTGGGGTCT
GTGCAGATCTGCTGGATCTAGGTCCAGGGAGTCTCAGTGATGGTCTGAGCCTGGCCGCGCCAGGCTGGG
TGTCAGAGAGCCAGGAATCGAAATGCTTCATGGAACAACCACCTGGCCTTCAAGTCCGCCATGGA
GTCATAGTTGAGCTGACTCCAGGGCTACAGCGGGTCTTACATTGCCTCCAGACGGTGAAGAAGGTGA
TAGAGATCAACCCATACCTGCTTGGCACCATGGCTGGGGCGCAGCGGATTGCAGCTTCTGGGAACGGCT
GTTGGCTCGGCAATGTCGAATCTATGAGCTTCGAAATAAGGAACGCATCTCTGTAGCAGCTGCCTCCAAA
CTGCTTGCCAACATGGTGTATCAGTACAAAGGCATGGGGCTGTCCATGGGCACCATGATCTGTGGCTGGG
ATAAGAGAGGCCCTGGCCTCTACTACGTGGACAGTGAAGGGAACCGGATTCAGGGGCCACCTTCTCTGT
AGGTTCTGGCTCTGTGTATGCATATGGGGTCTGGATCGGGGCTATTCTATGACCTGGAAGTGGAGCAG
GCCTATGATCTGGCCGTCGAGCCATCTACCAAGCCACCTACAGAGATGCCTACTCAGGAGGTGCAGTCA
ACCTCTACCACGTGCGGGAGGATGGCTGGATCCGAGTCTCCAGTGACAATGTGGCTGATCTACATGAGAA
GTATAGTGGCTTACCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



Protein Sequence: >RC209326 protein sequence
 Red=Cloning site Green=Tags(s)

MALASVLERPLPVNQRGFFGLGGRADLLDLGPGSLSDGLSLAAPGWGVPEEPGIEMLHGTTTLAFKFRHG
 VIVAADSRATAGAYIASQTVKKVIEINPYLLGTMAGGAADCSFWERLLARQCRIYELRNKERISVAAASK
 LLANMVYQYKGMGLSMGTMICGWDKRGPLYVDSEGNRISGATFSVSGSVYAYGVMDRGYSYDLEVEQ
 AYDLARRAIYQATYRDAYS GGAVNLYHVREDGWIRVSSDNVADLHEKYSGSTP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6126_d01.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_002797

ORF Size: 789 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_002797.5
RefSeq Size:	1311 bp
RefSeq ORF:	792 bp
Locus ID:	5693
UniProt ID:	P28074
Cytogenetics:	14q11.2
Domains:	proteasome
Protein Families:	Protease
Protein Pathways:	Proteasome
MW:	28.5 kDa
Gene Summary:	<p>The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit in the proteasome. This catalytic subunit is not present in the immunoproteasome and is replaced by catalytic subunit 3i (proteasome beta 8 subunit). Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2009]</p>


Product images:



Circular map for RC209326



Western blot validation of overexpression lysate (Cat# [LY400989]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC209326 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified PSMB5 protein (Cat# [TP309326]). The protein was produced from HEK293T cells transfected with PSMB5 cDNA clone (Cat# RC209326) using MegaTran 2.0 (Cat# [TT210002]).