

## Product datasheet for RC209392L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PRPS1L1 (NM\_175886) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** PRPS1L1 (NM\_175886) Human Tagged ORF Clone Lentiviral Particle

Symbol: PRPS1L1

Synonyms: PRPS1; PRPS3; PRPSL; PRS-III

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 175886

ORF Size: 954 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC209392).

Sequence:

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 175886.2

 RefSeq Size:
 1091 bp

 RefSeq ORF:
 957 bp

 Locus ID:
 221823

 UniProt ID:
 P21108

 Cytogenetics:
 7p21.1

**Protein Families:** Druggable Genome

**Protein Pathways:** Metabolic pathways, Pentose phosphate pathway, Purine metabolism





## PRPS1L1 (NM\_175886) Human Tagged ORF Clone Lentiviral Particle - RC209392L3V

**MW:** 34.8 kDa

**Gene Summary:** This intronless gene is specifically expressed in the testis, and encodes a protein that is highly

homologous to the two subunits of phosphoribosylpyrophosphate synthetase encoded by human X-linked genes, PRPS1 and PRPS2. These enzymes convert pyrimidine, purine or

pyridine bases to the corresponding ribonucleoside monophosphates. In vitro

transcription/translation and site-directed mutagenesis studies indicate that translation of this mRNA initiates exclusively at a non-AUG (ACG) codon. [provided by RefSeq, Jul 2008]