

Product datasheet for RC209451L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

VEGFC (NM_005429) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: VEGFC (NM 005429) Human Tagged ORF Clone Lentiviral Particle

Symbol: VEGFC

Synonyms: Flt4-L; LMPH1D; LMPHM4; VRP

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_005429 **ORF Size:** 1257 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209451).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 005429.2

 RefSeq Size:
 2103 bp

 RefSeq ORF:
 1260 bp

 Locus ID:
 7424

 UniProt ID:
 P49767

 Cytogenetics:
 4q34.3

Domains: PDGF, CXCXC

Protein Families: Druggable Genome, Secreted Protein





VEGFC (NM_005429) Human Tagged ORF Clone Lentiviral Particle - RC209451L2V

Protein Pathways: Bladder cancer, Cytokine-cytokine receptor interaction, Focal adhesion, mTOR signaling

pathway, Pancreatic cancer, Pathways in cancer, Renal cell carcinoma

MW: 46.9 kDa

Gene Summary: The protein encoded by this gene is a member of the platelet-derived growth factor/vascular

endothelial growth factor (PDGF/VEGF) family. The encoded protein promotes angiogenesis and endothelial cell growth, and can also affect the permeability of blood vessels. The proprotein is further cleaved into a fully processed form that can bind and activate VEGFR-2

and VEGFR-3 receptors. [provided by RefSeq, Apr 2014]