

Product datasheet for RC209462L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ATP6V1B1 (NM 001692) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ATP6V1B1 (NM_001692) Human Tagged ORF Clone Lentiviral Particle

Symbol:

ATP6B1; DRTA2; RTA1B; VATB; VMA2; VPP3 Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Myc-DDK Tag: NM 001692 ACCN: **ORF Size:** 1539 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC209462).

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer:

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001692.3

RefSeq Size: 1956 bp RefSeq ORF: 1542 bp

Locus ID: 525 **UniProt ID:** P15313 2p13.3 Cytogenetics:

Domains: ATP-synt_ab, ATP-synt_ab_C, ATP-synt_ab_N

Protein Families: Druggable Genome





ATP6V1B1 (NM_001692) Human Tagged ORF Clone Lentiviral Particle - RC209462L3V

Protein Pathways: Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative

phosphorylation, Vibrio cholerae infection

MW: 56.8 kDa

Gene Summary: This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that

mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is one of two V1 domain B subunit isoforms and is found in the kidney. Mutations in this gene cause distal renal tubular acidosis associated with sensorineural

deafness. [provided by RefSeq, Jul 2008]