

Product datasheet for RC209482L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Hexokinase II (HK2) (NM 000189) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Hexokinase II (HK2) (NM 000189) Human Tagged ORF Clone Lentiviral Particle

Symbol: Hexokinase II

HKII: HXK2 Synonyms:

Mammalian Cell

None

Selection: Vector:

pLenti-C-mGFP (PS100071)

mGFP Tag:

NM 000189 ACCN: **ORF Size:** 2751 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209482).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer:

> reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 000189.4

RefSeq Size: 7109 bp RefSeq ORF: 2754 bp Locus ID: 3099 **UniProt ID:** P52789 Cytogenetics: 2p12

Domains: hexokinase

Protein Families: Druggable Genome





Hexokinase II (HK2) (NM_000189) Human Tagged ORF Clone Lentiviral Particle - RC209482L2V

Protein Pathways: Amino sugar and nucleotide sugar metabolism, Fructose and mannose metabolism,

Galactose metabolism, Glycolysis / Gluconeogenesis, Insulin signaling pathway, Metabolic

pathways, Starch and sucrose metabolism, Type II diabetes mellitus

MW: 102.2 kDa

Gene Summary: Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most

glucose metabolism pathways. This gene encodes hexokinase 2, the predominant form found in skeletal muscle. It localizes to the outer membrane of mitochondria. Expression of this gene is insulin-responsive, and studies in rat suggest that it is involved in the increased rate

of glycolysis seen in rapidly growing cancer cells. [provided by RefSeq, Apr 2009]