

Product datasheet for RC209487L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Sec8 (EXOC4) (NM_021807) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Sec8 (EXOC4) (NM 021807) Human Tagged ORF Clone Lentiviral Particle

Symbol: Sec8

Synonyms: SEC8; SEC8L1; Sec8p

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_021807 **ORF Size:** 2922 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209487).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 021807.3

 RefSeq Size:
 4203 bp

 RefSeq ORF:
 2925 bp

 Locus ID:
 60412

 UniProt ID:
 Q96A65

 Cytogenetics:
 7q33

Domains: Sec8_exocyst

Protein Pathways: Tight junction





ORIGENE

MW: 110.5 kDa

Gene Summary:

The protein encoded by this gene is a component of the exocyst complex, a multiple protein complex essential for targeting exocytic vesicles to specific docking sites on the plasma membrane. Though best characterized in yeast, the component proteins and functions of exocyst complex have been demonstrated to be highly conserved in higher eukaryotes. At least eight components of the exocyst complex, including this protein, are found to interact with the actin cytoskeletal remodeling and vesicle transport machinery. The complex is also essential for the biogenesis of epithelial cell surface polarity. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]