

Product datasheet for RC209509L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SF3B3 (NM_012426) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SF3B3 (NM_012426) Human Tagged ORF Clone Lentiviral Particle

Symbol: SF3B3

Synonyms: RSE1; SAP130; SF3b130; STAF130

Mammalian Cell

None

Selection:

Vector:

pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_012426

ORF Size: 3651 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209509).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 012426.3

 RefSeq Size:
 9736 bp

 RefSeq ORF:
 3654 bp

 Locus ID:
 23450

 UniProt ID:
 Q15393

 Cytogenetics:
 16q22.1

 Domains:
 CPSF_A

Protein Pathways: Spliceosome





ORIGENE

MW: 135.5 kDa

Gene Summary: This gene encodes subunit 3 of the splicing factor 3b protein complex. Splicing factor 3b,

together with splicing factor 3a and a 12S RNA unit, forms the U2 small nuclear ribonucleoproteins complex (U2 snRNP). The splicing factor 3b/3a complex binds pre-mRNA upstream of the intron's branch site in a sequence independent manner and may anchor the U2 snRNP to the pre-mRNA. Splicing factor 3b is also a component of the minor U12-type spliceosome. Subunit 3 has also been identified as a component of the STAGA (SPT3-TAF(II)31-GCN5L acetylase) transcription coactivator-HAT (histone acetyltransferase) complex, and the TFTC (TATA-binding-protein-free TAF(II)-containing complex). These complexes may

function in chromatin modification, transcription, splicing, and DNA repair. [provided by

RefSeq, Jul 2008]