

Product datasheet for **RC209528**

ZCCHC8 (NM_017612) Human Tagged ORF Clone

Product data:

| | |
|---------------------------|-------------------------------------------|
| Product Type: | Expression Plasmids |
| Product Name: | ZCCHC8 (NM_017612) Human Tagged ORF Clone |
| Tag: | Myc-DDK |
| Symbol: | ZCCHC8 |
| Synonyms: | PFBMFT5 |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |



[View online »](#)

ORF Nucleotide
Sequence:

>RC209528 ORF sequence
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCCGCGATCGCC

ATGGCCGCAGAGGTGATTTTGGCGATCTAGAGCTCTTCGAGCCGTTTCGACCACCCAGAGGAGTCGATTC
CGAAGCCCGTTCCACTCGCTTCAAGGACGACGACGGCGACGAGGAGGACGAAAAATGGGGTCGGCGACGC
GGAGCTACGGGAGCGGCTTCGGCAGTGCAGGAGACCATCGAGCAGCTCCGCGCCGAGAATCAAGAAGCTT
AAACGAAAATTGAACATTCTGACTCGACCGAGTGAATATTGGTGAACGATACTAAGTTAGATGGACCTA
TATTACAGATTCTATTCAAGAATGCTATTTCAAAGCAATATCATCAAGAAATAGAGGAATTTGTATC
AAATTTAGTAAAAAGATTTGAGGAACAGCAGAAAAATGATGTGAAAAAGACTTCCCTTTAATCTTTTGCCC
CAGCCATCCAGTATTGTGCTAGAGGAGGACCACAAAGTGAAGAGTCTGTGCCATTAACAAACAACAGG
AAGCTTTCAGTGTGTAGGAAGTGTCTGTATTTACTAATTTTTGCCTTGATAAATGGGGCAACCGCT
TCTAAATGAAAACCTCAGCTTTCCGAAGGATGGGAAATACCCAAGTACCATCAAGTCTTCAGCCACATT
GTTTCTCTAGAAGGGCAAGAAATACAAGTAAAGGCAAAAAGGCCAAAGCCTCACTGTTTCAATTGTGGTT
CTGAAGAACACCAATGAAGATTGCCAATGCCTCGGAATGCTGCTCGAATAAGTGAAGAGAGAAAAGA
GTATATGGATGCCTGTGGAGAAGCAACAATCAGAATTTCCAGCAGCGATACCACGCAGAAGAGTAGAA
GAAAGATTTGGAAGATTCAGCCAGGAGTTATTAGTGAGGAAGTCAAGATGCCTAGGTGTGACAGACA
AGAGTCTTCCACCTTTTATCTATCGGATGCGCCAGCTAGGGTACCCACCAGGGTGGCTCAAAGAGGTTGA
ATTGGAGAATTCGGGGCTTGACTCTATGATGAAAAAGATGGCACTGATGGGAAACAGAAGTTGGAGAA
ATACAACAGAATAAAGTGTCACTTACGATCTCTCAAAATTTGGTCAACTATCCTGGTTTTAATATATCTA
CTCCCAGAGGAATCCAGACGAATGGAGGATCTTTGGTTCCATACCAATGCAGGCATGCAGCAGAAGGA
TGTGTTTGCCAATTACCTTACTTCTAACTTCCAAGCGCCAGGTGTGAAGTCTGGCAACAAGAGGTCTTCA
TCTCACTCTAGCCAGGTAGTCCAAGAAGCAGAAGAATGAAAGCAACTCAGCGGGATCTCCCGCCGACA
TGGAGCTCGATTAGATATGGAGGTACCACATGGTTCTCAGAGCAGCGAAAGTTTTTCAGTTTCAACCACC
ATTACCTCTGACTCTCCACTCCCCCGGGAACTCTCCACCCGTCTTACCCTCCACTCCCAAAG
GGCACCCCGCGCTGACTCCCAGTACTACCCAGACCAGAACAGCATCTGGAGCTGTGGATGAGGACG
CACTGACTCTAGAAGAACTTGAAGAACAGCAGAGGCGGATCTGGGAGCTCTTGGCAGGCGGAGAGCGT
AAACAGCGACTCCGACGTTCTGTGGACACACCTTTAACTGGCAATTCGGTTGCCTCATCACCTTGTCCA
AATGAGCTAGACCTCCCTGTCCCGGAGGAAAAACATCTGAAAAGCAGACGCTGGATGAGCCTGAGGTAC
CAGAGATTTTTACAAAGAAATCAGAAGCTGGACATGCCTCCAGTCCAGACTCTGAGGTGACATCACTTTG
TCAGAAGGAAAAAGCAGAGTTGGCTCCGGTAAACTGAAAGTGCCTTCTTGATAATGGCAGTGTGTA
CCAAACTGTGACATCAGCAATGGGGCAGCCAGAAGCTCTTTCCTGCAGACACCAGTCTTCAACGGCCA
CTAAAATTCATAGCCCTATACCTGACATGAGCAAAATTTGCAACTGGAATCACGCCATTTGAATTTGAGAA
TATGGCAGAACTACTGGAATGTACCTCAGGATAAGAAGCTTGTAAAGAAGTCAACCCGAAACCAGCAG
AAAAACAAAAGGCTCTGAA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC209528 protein sequence
Red=Cloning site Green=Tags(s)

MAAEVYFGDLELFEFPDHPEESIPKPVHTRFKDDDDGDEEDENGVGDAELRERLRQCEETIEQLRAENQEL
 KRKLNILTRPSGILVNDTKLDGPILQILFMNNAISKQYHQEIEEFVSNL VKRFEEQKNDVEKTSFNLLP
 QPSSIVLEEDHKVEESCAIKNNKEAFSVVGSVL YFTNFCLDKLGQPLL NENPQLSEGWEIPKYHQVFSHI
 VSLEGQEIQVKAKRKPFCNCGSEEHQMKDCMPRNaARI SEKRKEYMDACGEANNQNFQORYHAEEVE
 ERFGRFRKPGVISEELQDALGVTDKSLPPFIYRMRQLGYPPGWLKEVELENSGLALYDGKDGTDGETEVE
 IQQNKSVTYDLSKLVNYPGFNISTPRGIPDEWRIFGSIPMQACQKQDVFANYLTSNFQAPGVKSGNKRSS
 SHSSPGSPKKQKNESNSAGSPADMELDSMEVPHGSQSSESFQFQPPLPDPPLPRGTPPPVFTPLPK
 GTPPLTPSDSPQTRTASGAVDEDAL TLEELQQRRIWAALQAEVNSDSDVPVDTPLTGNSVASSPCP
 NELDLPVPEGKTEKQTLDEPEVPEIFTKSEAGHASSPDSEVTS LCQKEKAELAPVNTGALLDNGSVV
 PNCDISNGGSQKLFADTSPSTATKIHSPIPDMSKFATGITPFEFENMAESTGMYLRIRSLKNSPRNQ
 KNKKASE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6356_f06.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_017612


ORF Size: 2121 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)


OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

| | |
|-------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | <ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | <u>NM_017612.5</u> |
| RefSeq Size: | 4260 bp |
| RefSeq ORF: | 2124 bp |
| Locus ID: | 55596 |
| UniProt ID: | <u>Q6NZY4</u> |
| Cytogenetics: | 12q24.31 |
| Domains: | PSP |
| MW: | 78.6 kDa |
| Gene Summary: | This gene encodes a scaffold protein which serves as an assessory factor to the nuclear RNA exosome complex. The encoded protein forms a trimeric human nuclear exosome targeting (NEXT) complex, together with hMTR4 and the RNA-binding factor RBM7 which promotes the exosomal degradation of non-coding promoter-upstream transcripts, enhancer RNAs and 3'-extended products of histone- and small nuclear RNA transcription. This complex is also thought to recruit the exosome to degrade intronic RNAs via its interaction with both the exosome and the spliceosome. It contains both an N-terminal zinc-knuckle domain and a C-terminal proline-rich domain. [provided by RefSeq, Apr 2017] |


Product images:



Circular map for RC209528



Western blot validation of overexpression lysate (Cat# [LY413673]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC209528 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified ZCCHC8 protein (Cat# [TP309528]). The protein was produced from HEK293T cells transfected with ZCCHC8 cDNA clone (Cat# RC209528) using MegaTran 2.0 (Cat# [TT210002]).