

## Product datasheet for RC209627L4

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## ATG4D (NM\_032885) Human Tagged Lenti ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: ATG4D (NM\_032885) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: ATG4D

**Synonyms:** APG4-D; APG4D; AUTL4

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC209627).

Sequence:

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_032885

ORF Size: 1422 bp



#### ATG4D (NM\_032885) Human Tagged Lenti ORF Clone - RC209627L4

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Regulation of autophagy

**RefSeq:** NM 032885.4

 RefSeq Size:
 1929 bp

 RefSeq ORF:
 1425 bp

 Locus ID:
 84971

 UniProt ID:
 Q86TL0

Cytogenetics: 19p13.2

**Domains:** Peptidase\_C54

**MW:** 52.7 kDa

**Protein Pathways:** 

**Gene Summary:** Autophagy is the process by which endogenous proteins and damaged organelles are


destroyed intracellularly. Autophagy is postulated to be essential for cell homeostasis and cell remodeling during differentiation, metamorphosis, non-apoptotic cell death, and aging. Reduced levels of autophagy have been described in some malignant tumors, and a role for autophagy in controlling the unregulated cell growth linked to cancer has been proposed. This gene belongs to the autophagy-related protein 4 (Atg4) family of C54 endopeptidases.

Members of this family encode proteins that play a role in the biogenesis of

autophagosomes, which sequester the cytosol and organelles for degradation by lysosomes. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013]



# **Product images:**



Circular map for RC209627L4