

## Product datasheet for RC209676L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## C5L2 (C5AR2) (NM\_018485) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** C5L2 (C5AR2) (NM\_018485) Human Tagged ORF Clone Lentiviral Particle

Symbol: C5L2

**Synonyms:** C5L2; GPF77; GPR77

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 018485

ORF Size: 1011 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC209676).

Sequence:

**Domains:** 

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 018485.1

 RefSeq Size:
 1396 bp

 RefSeq ORF:
 1014 bp

 Locus ID:
 27202

 UniProt ID:
 Q9P296

 Cytogenetics:
 19q13.32

\_

**Protein Families:** Druggable Genome, GPCR, Transmembrane

7tm 1





ORIGENE

**MW:** 36.1 kDa

**Gene Summary:** This gene encodes a G-protein coupled receptor 1 family member involved in the

complement system of the innate immune response. Unlike classical G-protein coupled receptors, the encoded protein does not associate with intracellular G-proteins. It may instead modulate signal transduction through the beta-arrestin pathway, and may alternatively act as a decoy receptor. This gene may be involved in coronary artery disease and in the pathogenesis of sepsis. Alternative splicing results in multiple transcript variants.

[provided by RefSeq, Nov 2012]