

# Product datasheet for RC209687L3

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL7R alpha (IL7R) (NM\_002185) Human Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

Product Name: IL7R alpha (IL7R) (NM\_002185) Human Tagged Lenti ORF Clone

Tag: Myc-DDK
Symbol: IL7R alpha

Synonyms: CD127; CDW127; IL-7R-alpha; IL7RA; ILRA

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Sgfl-Mlul

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC209687).


Sequence:

Restriction Sites: Cloning Scheme:

Cloning sites used for ORF Shuttling:

Sgf 1 ORF Mlu I

---- GCG ATC GCC ATG ---//--- NNN ACG CGT ---



<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_002185

ORF Size: 1377 bp



## IL7R alpha (IL7R) (NM\_002185) Human Tagged Lenti ORF Clone - RC209687L3

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 002185.2</u>

 RefSeq Size:
 4617 bp

 RefSeq ORF:
 1380 bp

 Locus ID:
 3575

 UniProt ID:
 P16871

**Cytogenetics:** 5p13.2

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT signaling

pathway, Primary immunodeficiency

**MW:** 51.6 kDa


**Gene Summary:** The protein encoded by this gene is a receptor for interleukin 7 (IL7). The function of this

receptor requires the interleukin 2 receptor, gamma chain (IL2RG), which is a common gamma chain shared by the receptors of various cytokines, including interleukins 2, 4, 7, 9, and 15. This protein has been shown to play a critical role in V(D)J recombination during lymphocyte development. Defects in this gene may be associated with severe combined immunodeficiency (SCID). Alternatively spliced transcript variants have been found. [provided

by RefSeq, Dec 2015]



# **Product images:**



Circular map for RC209687L3