

Product datasheet for RC209796L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TIMP2 (NM_003255) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TIMP2 (NM_003255) Human Tagged ORF Clone Lentiviral Particle

Symbol: TIMP2

Synonyms: CSC-21K; DDC8

Mammalian Cell

Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_003255

ORF Size: 660 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209796).

Sequence:

Domains:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 003255.4</u>

 RefSeq Size:
 3670 bp

 RefSeq ORF:
 663 bp

 Locus ID:
 7077

 UniProt ID:
 P16035

Cytogenetics: 17q25.3

Protein Families: Druggable Genome, Secreted Protein

NTR



ORIGENE

MW: 24.4 kDa

Gene Summary:

This gene is a member of the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix. In addition to an inhibitory role against metalloproteinases, the encoded protein has a unique role among TIMP family members in its ability to directly suppress the proliferation of endothelial cells. As a result, the encoded protein may be critical to the maintenance of tissue homeostasis by suppressing the proliferation of quiescent tissues in response to angiogenic factors, and by inhibiting protease activity in tissues undergoing remodelling of the extracellular matrix. [provided by RefSeq, Jul 2008]