

## Product datasheet for RC209810L3

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# C13orf15 (RGCC) (NM\_014059) Human Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** C13orf15 (RGCC) (NM\_014059) Human Tagged Lenti ORF Clone

Tag: Myc-DDK
Symbol: C13orf15

Synonyms: bA157L14.2; C13orf15; RGC-32; RGC32

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC209810).

Sequence:

Restriction Sites: Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_014059

ORF Size: 411 bp





## C13orf15 (RGCC) (NM\_014059) Human Tagged Lenti ORF Clone - RC209810L3

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 014059.2</u>, <u>NP 054778.2</u>

 RefSeq Size:
 1126 bp

 RefSeq ORF:
 414 bp

 Locus ID:
 28984

 UniProt ID:
 Q9H4X1

 Cytogenetics:
 13q14.11

**MW:** 14.6 kDa


**Gene Summary:** This gene is thought to regulate cell cycle progression. It is induced by p53 in response to

DNA damage, or by sublytic levels of complement system proteins that result in activation of the cell cycle. The encoded protein localizes to the cytoplasm during interphase and to centrosomes during mitosis. The protein forms a complex with polo-like kinase 1. The protein also translocates to the nucleus in response to treatment with complement system proteins, and can associate with and increase the kinase activity of cell division cycle 2 protein. In different assays and cell types, overexpression of this protein has been shown to activate or

suppress cell cycle progression. [provided by RefSeq, Jul 2008]



# **Product images:**



Circular map for RC209810L3