

## Product datasheet for RC209837L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RHOB (NM 004040) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** RHOB (NM\_004040) Human Tagged ORF Clone Lentiviral Particle

Symbol:

ARH6; ARHB; MST081; MSTP081; RHOH6 Synonyms:

**Mammalian Cell** 

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 004040 ACCN:

**ORF Size:** 588 bp

**ORF Nucleotide** 

Sequence: OTI Disclaimer: The ORF insert of this clone is exactly the same as(RC209837).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 004040.2

RefSeq Size: 2384 bp RefSeq ORF: 591 bp Locus ID: 388 **UniProt ID:** P62745

Cytogenetics: 2p24.1 **Domains:** 

**Protein Families:** Druggable Genome

ras, RAS, RHO, RAB







**MW:** 22.1 kDa

**Gene Summary:** 

Mediates apoptosis in neoplastically transformed cells after DNA damage. Not essential for development but affects cell adhesion and growth factor signaling in transformed cells. Plays a negative role in tumorigenesis as deletion causes tumor formation. Involved in intracellular protein trafficking of a number of proteins. Targets PKN1 to endosomes and is involved in trafficking of the EGF receptor from late endosomes to lysosomes. Also required for stability and nuclear trafficking of AKT1/AKT which promotes endothelial cell survival during vascular development. Serves as a microtubule-dependent signal that is required for the myosin contractile ring formation during cell cycle cytokinesis. Required for genotoxic stress-induced cell death in breast cancer cells.[UniProtKB/Swiss-Prot Function]