

### **Product datasheet for RC209842**

# SC35 (SRSF2) (NM 003016) Human Tagged ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** SC35 (SRSF2) (NM\_003016) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: SC35

Synonyms: PR264; SC-35; SC35; SFRS2; SFRS2A; SRp30b

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >RC209842 representing NM\_003016

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

 ${\tt TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC}$ 

GCCGCGATCGCC

AAGTCTCCTGAAGAGGAAGGAGCGGTGTCCTCT

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >RC209842 representing NM\_003016

Red=Cloning site Green=Tags(s)


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mg4841">https://cdn.origene.com/chromatograms/mg4841</a> h09.zip

**Restriction Sites:** Sgfl-Mlul

Cloning Scheme:





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_003016

ORF Size: 663 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customercom">customercom</a> or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



Domains:

#### SC35 (SRSF2) (NM\_003016) Human Tagged ORF Clone - RC209842

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 003016.3</u>

 RefSeq Size:
 2923 bp

 RefSeq ORF:
 666 bp

 Locus ID:
 6427

 UniProt ID:
 Q01130

 Cytogenetics:
 17q25.1

**Protein Families:** Stem cell - Pluripotency, Transcription Factors

RRM

Protein Pathways: Spliceosome MW: 25.3 kDa

**Gene Summary:** The protein encoded by this gene is a member of the serine/arginine (SR)-rich family of pre-


mRNA splicing factors, which constitute part of the spliceosome. Each of these factors contains an RNA recognition motif (RRM) for binding RNA and an RS domain for binding other proteins. The RS domain is rich in serine and arginine residues and facilitates interaction between different SR splicing factors. In addition to being critical for mRNA splicing, the SR proteins have also been shown to be involved in mRNA export from the nucleus and in translation. Two transcript variants encoding the same protein and one non-coding transcript

variant have been found for this gene. In addition, a pseudogene of this gene has been found


on chromosome 11. [provided by RefSeq, Sep 2010]



## **Product images:**



Circular map for RC209842



Western blot validation of overexpression lysate (Cat# [LY418954]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC209842 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).