

Product datasheet for RC209920L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HLA (HLA-DRA) (NM_019111) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: HLA (HLA-DRA) (NM_019111) Human Tagged ORF Clone Lentiviral Particle

Symbol: HLA-DRA
Synonyms: HLA-DRA1

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_019111

ORF Size: 762 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC209920).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 019111.3

 RefSeq Size:
 1312 bp

 RefSeq ORF:
 765 bp

 Locus ID:
 3122

 UniProt ID:
 P01903

 Cytogenetics:
 6p21.32

Domains: MHC_II_alpha, ig, IGc1

Protein Families: Transmembrane





HLA (HLA-DRA) (NM_019111) Human Tagged ORF Clone Lentiviral Particle - RC209920L1V

Protein Pathways: Allograft rejection, Antigen processing and presentation, Asthma, Autoimmune thyroid

disease, Cell adhesion molecules (CAMs), Graft-versus-host disease, Hematopoietic cell

lineage, Systemic lupus erythematosus, Type I diabetes mellitus, Viral myocarditis

MW: 28.6 kDa

Gene Summary: HLA-DRA is one of the HLA class II alpha chain paralogues. This class II molecule is a

heterodimer consisting of an alpha and a beta chain, both anchored in the membrane. This

molecule is expressed on the surface of various antigen presenting cells such as B

lymphocytes, dendritic cells, and monocytes/macrophages, and plays a central role in the immune system and response by presenting peptides derived from extracellular proteins, in particular, pathogen-derived peptides to T cells. The alpha chain is approximately 33-35 kDa and its gene contains 5 exons. Exon 1 encodes the leader peptide, exons 2 and 3 encode the

two extracellular domains, and exon 4 encodes the transmembrane domain and the cytoplasmic tail. DRA does not have polymorphisms in the peptide binding part and acts as

the sole alpha chain for DRB1, DRB3, DRB4 and DRB5. [provided by RefSeq, Aug 2020]