

Product datasheet for RC209946L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GGT5 (NM_004121) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: GGT5 (NM_004121) Human Tagged ORF Clone Lentiviral Particle

Symbol:

GGL; GGT-REL; GGT 5; GGTLA1 Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 004121 ACCN: **ORF Size:** 1761 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC209946).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 004121.1

RefSeq Size: 2497 bp RefSeq ORF: 1761 bp Locus ID: 2687 **UniProt ID:** P36269 Cytogenetics: 22q11.23

Domains: G_glu_transpept

Protein Families: Protease, Transmembrane





GGT5 (NM_004121) Human Tagged ORF Clone Lentiviral Particle - RC209946L3V

Protein Pathways: Arachidonic acid metabolism, Cyanoamino acid metabolism, Glutathione metabolism,

Metabolic pathways, Selenoamino acid metabolism, Taurine and hypotaurine metabolism

MW: 62.3 kDa

Gene Summary: This gene is a member of the gamma-glutamyl transpeptidase gene family, and some reports

indicate that it is capable of cleaving the gamma-glutamyl moiety of glutathione. The protein encoded by this gene is synthesized as a single, catalytically-inactive polypeptide, that is processed post-transcriptionally to form a heavy and light subunit, with the catalytic activity contained within the small subunit. The encoded enzyme is able to convert leukotriene C4 to leukotriene D4, but appears to have distinct substrate specificity compared to gamma-glutamyl transpeptidase. Alternative splicing results in multiple transcript variants encoding

different isoforms. [provided by RefSeq, Oct 2014]