

Product datasheet for RC209949L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

VDAC1 (NM_003374) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: VDAC1 (NM_003374) Human Tagged ORF Clone Lentiviral Particle

Symbol: VDAC1

Synonyms: PORIN; VDAC-1

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_003374

ORF Size: 849 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209949).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 003374.1</u>

RefSeq Size: 1806 bp
RefSeq ORF: 852 bp
Locus ID: 7416
UniProt ID: P21796

Cytogenetics: 5q31.1

Domains: Euk_porin

Protein Families: Ion Channels: Other





VDAC1 (NM_003374) Human Tagged ORF Clone Lentiviral Particle - RC209949L2V

Protein Pathways: Calcium signaling pathway, Huntington's disease, Parkinson's disease

MW: 30.6 kDa

Gene Summary: This gene encodes a voltage-dependent anion channel protein that is a major component of

the outer mitochondrial membrane. The encoded protein facilitates the exchange of metabolites and ions across the outer mitochondrial membrane and may regulate

mitochondrial functions. This protein also forms channels in the plasma membrane and may

be involved in transmembrane electron transport. Alternate splicing results in multiple transcript variants. Multiple pseudogenes of this gene are found on chromosomes 1, 2 3, 6, 9,

12, X and Y.[provided by RefSeq, Sep 2010]