

## Product datasheet for RC209951L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IRF3 (NM\_001571) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** IRF3 (NM\_001571) Human Tagged ORF Clone Lentiviral Particle

Symbol: IRF3
Synonyms: IIAE7

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_001571

 ORF Size:
 1281 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC209951).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001571.2</u>

 RefSeq Size:
 1626 bp

 RefSeq ORF:
 1284 bp

 Locus ID:
 3661

 UniProt ID:
 Q14653

 Cytogenetics:
 19q13.33

Domains: IRF

**Protein Families:** Druggable Genome, Transcription Factors





## IRF3 (NM\_001571) Human Tagged ORF Clone Lentiviral Particle - RC209951L1V

**Protein Pathways:** Cytosolic DNA-sensing pathway, RIG-I-like receptor signaling pathway, Toll-like receptor

signaling pathway

**MW:** 47.2 kDa

**Gene Summary:** This gene encodes a member of the interferon regulatory transcription factor (IRF) family.

The encoded protein is found in an inactive cytoplasmic form that upon serine/threonine phosphorylation forms a complex with CREBBP. This complex translocates to the nucleus and activates the transcription of interferons alpha and beta, as well as other interferon-induced genes. The protein plays an important role in the innate immune response against DNA and RNA viruses. Mutations in this gene are associated with Encephalopathy, acute, infection-

induced, herpes-specific, 7. [provided by RefSeq, Sep 2020]