

## Product datasheet for RC209966L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Retinoic Acid Receptor alpha (RARA) (NM\_001024809) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Retinoic Acid Receptor alpha (RARA) (NM\_001024809) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Retinoic Acid Receptor alpha

Synonyms: NR1B1; RAR

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM 001024809

ORF Size: 1371 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC209966).

Sequence:
OTI Disclaimer:

Cytogenetics:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 001024809.2

17q21.2

 RefSeq Size:
 3494 bp

 RefSeq ORF:
 1374 bp

 Locus ID:
 5914

 UniProt ID:
 P10276

**Protein Families:** Druggable Genome, Nuclear Hormone Receptor, Transcription Factors





## Retinoic Acid Receptor alpha (RARA) (NM\_001024809) Human Tagged ORF Clone Lentiviral Particle – RC209966L2V

**Protein Pathways:** Acute myeloid leukemia, Pathways in cancer

MW: 50.7 kDa

**Gene Summary:** This gene represents a nuclear retinoic acid receptor. The encoded protein, retinoic acid

receptor alpha, regulates transcription in a ligand-dependent manner. This gene has been implicated in regulation of development, differentiation, apoptosis, granulopoeisis, and transcription of clock genes. Translocations between this locus and several other loci have been associated with acute promyelocytic leukemia. Alternatively spliced transcript variants

have been found for this locus.[provided by RefSeq, Sep 2010]