

Product datasheet for RC209977

LIPT1 (NM_145198) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	LIPT1 (NM_145198) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	LIPT1
Synonyms:	LIPT1D
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC209977 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGCTGATCCCATTTTCAATGAAGAATTGCTTCCAGTTACTTTGTAAGTCCAGGTTCCAGCAGCTGGCT
TAAAAAACAGTAAAAATGGGCTCATTTTACAGTCAATTTCCAATGATGTCTATCAAAATCTGGCTGT
GGAAGACTGGATCCATGACCATATGAATCTAGAAGGCAAACCAATTCTATTCTTTGGCAGAATTCTCC
TCTGTTGTAATTGGTAGGCATCAAAATCCTTGGCAGGAATGTAACCTGAATCTAATGAGAGAAGAAGGTA
TAAACTGGCTCGGAGAAGAAGTGGAGGAGGAACAGTCTACCATGATATGGGTAATATCAATTTGACTTT
CTTTACAACCAAAAAAGTATGATAGAATGGAAAATCTGAAATTAATTGTGAGAGCTCTGAATGCTGTC
CAACCCAGCTGGATGTGCAGGCTACCAAAAGATTTGACCTTTTACTTGATGGACAGTTTAAATCTCAG
GAACAGCTTCTAAGATCGGCCGGACTACTGCCTATCACCATTGCACCTTTATTATGTAGTACTGATGGGAC
GTTCTTGTCTTTGCTAAAGAGCCCTTACCAAGGATCAGGAGCAATGCCACTGCTAGCATACCTTCC
TTAGTGAAAAATCTTTGGAAAAGGATCCCACTCTGACCTGTGAAGTACTAATGAATGCTGTTGCTACAG
AGTATGCTGCTTATCATCAAATGATAATCACATTCACCTAATAAACCAACGGATGAGACACTGTTTCC
TGAATAAATAGCAAAGCCAAAGAAGTCAAACCTGGGAGTGGATATATGGCAAACTCCAAAGTTAGT
ATAAATACTTCTTTTATGTTATATGAACAGTCACACTTGGAAATTAAGTATTCATAGACATAAAGA
ATGGAAGAATTGAAATTTGTAATTGAAACACCTGATCATTGGTTGCCATTGGAAATACGTGACAAATT
AAATTCAGTCTTATTGGCAGTAAGTTTGGCCAACTGAAACTACCATGCTAACAAATATATTACTTAGA
ACATGTCCACAAGACCACAACTAAACAGTAAATGGAATATTCTCTGTGAAAAAATTAAGGAATAATG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC209977 protein sequence
Red=Cloning site Green=Tags(s)

MLIPFSMKNCFQLLCNCQVPAAGFKKTVKNGLI LQSI SNDVYQNLAVEDWIHDHNMNLEGPILFFWQNSP
 SVVIGRHQNPWQECNLNLMREEGIKLARRRSGGGTVYHDMGNINL TFFT KKKYDRMENLKLIVRALNAV
 QPQLDVQATKRFDLLLDGQFKISGTASKIGRTTAYHHCTLLCSTDGTFLSLLKSPYQGIRSNATASIPS
 LVKNLLEKDPTLTCEVLMNAVATEYAAYHQIDNHIHLINPTDETLPFGINSKAKELQTWEWIYGKTPKFS
 INTSFHVLVEQSHLEIKVFIDIKNGRIEICNIEAPDHWLPLEIRDKLNSSLIGSKFCPTETTMLTNILLR
 TCPQDHKLNKWNILCEKIKGIM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6557_g10.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_145198

ORF Size: 1119 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_145198.3](#)

RefSeq Size: 1572 bp

RefSeq ORF: 1122 bp

Locus ID: 51601

UniProt ID: [Q9Y234](#)


Cytogenetics: 2q11.2

Protein Pathways: Lipoic acid metabolism, Metabolic pathways


MW: 42.5 kDa

Gene Summary: The process of transferring lipoic acid to proteins is a two-step process. The first step is the activation of lipoic acid by lipoate-activating enzyme to form lipoyl-AMP. For the second step, the protein encoded by this gene transfers the lipoyl moiety to apoproteins. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 13. Read-through transcription also exists between this gene and the neighboring downstream mitochondrial ribosomal protein L30 (MRPL30) gene. [provided by RefSeq, Mar 2011]

Product images:



Circular map for RC209977



Western blot validation of overexpression lysate (Cat# [LY408046]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC218958] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).