

## **Product datasheet for RC209986**

## RPL32 (NM 000994) Human Tagged ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

Product Name: RPL32 (NM\_000994) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: RPL32

Synonyms: L32; PP9932

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC209986 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAG**GTTTAA** 

Protein Sequence: >RC209986 protein sequence

Red=Cloning site Green=Tags(s)

 ${\tt MAALRPLVKPKIVKKRTKKFIRHQSDRYVKIKRNWRKPRGIDNRVRRRFKGQILMPNIGYGSNKKTKHMLPSGFRKFLVHNVKELEVLLMCNKSYCAEIAHNVSSKNRKAIVERAAQLAIRVTNPNARLRSEENE}$ 

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mk6547">https://cdn.origene.com/chromatograms/mk6547</a> d06.zip

**Restriction Sites:** Sgfl-Mlul



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



#### **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_000994

ORF Size: 405 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** NM 000994.4

RefSeq Size: 1668 bp
RefSeq ORF: 408 bp
Locus ID: 6161
UniProt ID: P62910



Cytogenetics: 3p25.2

**Domains:** Ribosomal\_L32e

**Protein Pathways:** Ribosome

**MW:** 15.9 kDa


**Gene Summary:** Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and

a large 60S subunit. Together these subunits are composed of 4 RNA species and


approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L32E family of ribosomal proteins. It is located in the cytoplasm. Although some studies have mapped this gene to 3q13.3-q21, it is believed to map to 3p25-p24. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. Alternatively spliced transcript variants encoding the same protein have been

observed for this gene. [provided by RefSeq, Jul 2008]

# **Product images:**



Circular map for RC209986



Western blot validation of overexpression lysate (Cat# [LY424406]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC209986 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).