

## Product datasheet for RC209990L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RPL13A (NM\_012423) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: RPL13A (NM\_012423) Human Tagged ORF Clone Lentiviral Particle

Symbol: RPL13A

Synonyms: L13A; TSTA1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_012423

ORF Size: 609 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC209990).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 012423.2

 RefSeq Size:
 1196 bp

 RefSeq ORF:
 612 bp

 Locus ID:
 23521

 UniProt ID:
 P40429

 Cytogenetics:
 19q13.33

**Domains:** Ribosomal L13

**Protein Pathways:** Ribosome





ORIGENE

**MW:** 23.6 kDa

**Gene Summary:** 

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a member of the L13P family of ribosomal proteins that is a component of the 60S subunit. The encoded protein also plays a role in the repression of inflammatory genes as a component of the IFN-gamma-activated inhibitor of translation (GAIT) complex. This gene is co-transcribed with the small nucleolar RNA genes U32, U33, U34, and U35, which are located in the second, fourth, fifth, and sixth introns, respectively. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed throughout the genome. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jul 2012]