

Product datasheet for RC210043L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PPM1F (NM_014634) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PPM1F (NM_014634) Human Tagged ORF Clone Lentiviral Particle

Symbol: PPM1F

Synonyms: CAMKP; CaMKPase; FEM-2; hFEM-2; POPX2

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_014634 **ORF Size:** 3534 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC210043).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 014634.2</u>

 RefSeq Size:
 5199 bp

 RefSeq ORF:
 1365 bp

 Locus ID:
 9647

 UniProt ID:
 P49593

Cytogenetics: 22q11.22

Domains: PP2C

Protein Families: Druggable Genome, Phosphatase





ORIGENE

MW: 49.7 kDa

Gene Summary:

The protein encoded by this gene is a member of the PP2C family of Ser/Thr protein phosphatases. PP2C family members are known to be negative regulators of cell stress response pathways. This phosphatase can interact with Rho guanine nucleotide exchange factors (PIX), and thus block the effects of p21-activated kinase 1 (PAK), a protein kinase mediating biological effects downstream of Rho GTPases. Calcium/calmodulin-dependent protein kinase II gamma (CAMK2G/CAMK-II) is found to be one of the substrates of this phosphatase. The overexpression of this phosphatase or CAMK2G has been shown to mediate caspase-dependent apoptosis. An alternatively spliced transcript variant has been identified, but its full-length nature has not been determined. [provided by RefSeq, Jul 2008]