

Product datasheet for **RC210046L1V**

PSMD4 (NM_002810) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	PSMD4 (NM_002810) Human Tagged ORF Clone Lentiviral Particle
Symbol:	PSMD4
Synonyms:	AF; AF-1; ASF; MCB1; pUB-R5; Rpn10; S5A
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_002810
ORF Size:	1131 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC210046).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_002810.2
RefSeq Size:	1332 bp
RefSeq ORF:	1134 bp
Locus ID:	5710
UniProt ID:	P55036
Cytogenetics:	1q21.3
Domains:	VWA, UIM
Protein Pathways:	Proteasome



[View online »](#)

MW: 40.7 kDa

Gene Summary: The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the non-ATPase subunits of the 19S regulator lid. Pseudogenes have been identified on chromosomes 10 and 21. [provided by RefSeq, Jul 2008]