

## Product datasheet for RC210075L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RPS8 (NM\_001012) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type: Lentiviral Particles

**Product Name:** RPS8 (NM\_001012) Human Tagged ORF Clone Lentiviral Particle

Symbol: RPS8
Synonyms: S8

Mammalian Cell Puromycin

Selection:

i di Oiliyeiii

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_001012

ORF Size: 624 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC210075).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001012.1

 RefSeq Size:
 705 bp

 RefSeq ORF:
 627 bp

 Locus ID:
 6202

 UniProt ID:
 P62241

 Cytogenetics:
 1p34.1

**Domains:** Ribosomal\_S8e

**Protein Pathways:** Ribosome







**MW:** 24 kDa

**Gene Summary:** 

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 40S subunit. The protein belongs to the S8E family of ribosomal proteins. It is located in the cytoplasm. Increased expression of this gene in colorectal tumors and colon polyps compared to matched normal colonic mucosa has been observed. This gene is co-transcribed with the small nucleolar RNA genes U38A, U38B, U39, and U40, which are located in its fourth, fifth, first, and second introns, respectively. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008]