

## Product datasheet for **RC210081L2V**

### **MNK2 (MKNK2) (NM\_017572) Human Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	MNK2 (MKNK2) (NM_017572) Human Tagged ORF Clone Lentiviral Particle
Symbol:	MNK2
Synonyms:	GPRK7; MNK2
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_017572
ORF Size:	1242 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC210081).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_017572.3</a>
RefSeq Size:	1778 bp
RefSeq ORF:	1245 bp
Locus ID:	2872
UniProt ID:	<a href="#">Q9HBH9</a>
Cytogenetics:	19p13.3
Domains:	pkinese, TyrKc, S_TKc
Protein Families:	Druggable Genome, Protein Kinase



[View online »](#)

**Protein Pathways:** Insulin signaling pathway, MAPK signaling pathway

**MW:** 46.5 kDa

**Gene Summary:** This gene encodes a member of the calcium/calmodulin-dependent protein kinases (CAMK) Ser/Thr protein kinase family, which belongs to the protein kinase superfamily. This protein contains conserved DLG (asp-leu-gly) and ENIL (glu-asn-ile-leu) motifs, and an N-terminal polybasic region which binds importin A and the translation factor scaffold protein eukaryotic initiation factor 4G (eIF4G). This protein is one of the downstream kinases activated by mitogen-activated protein (MAP) kinases. It phosphorylates the eukaryotic initiation factor 4E (eIF4E), thus playing important roles in the initiation of mRNA translation, oncogenic transformation and malignant cell proliferation. In addition to eIF4E, this protein also interacts with von Hippel-Lindau tumor suppressor (VHL), ring-box 1 (Rbx1) and Cullin2 (Cul2), which are all components of the CBC(VHL) ubiquitin ligase E3 complex. Multiple alternatively spliced transcript variants have been found, but the full-length nature and biological activity of only two variants are determined. These two variants encode distinct isoforms which differ in activity and regulation, and in subcellular localization. [provided by RefSeq, Aug 2011]