

Product datasheet for RC210120

FGF10 (NM_004465) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
 Product Name: FGF10 (NM_004465) Human Tagged ORF Clone
 Tag: Myc-DDK
 Symbol: FGF10
 Mammalian Cell Selection: Neomycin
 Vector: pCMV6-Entry (PS100001)
 E. coli Selection: Kanamycin (25 ug/mL)
 ORF Nucleotide Sequence: >RC210120 representing NM_004465
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGTGGAATGGATACTGACACATTGTGCCTCAGCCTTCCCCACCTGCCGGCTGCTGCTGCTGCTGCT
 TTTTGTGCTGTTCTTGGTGTCTCCGTCCTGTACCTGCCAAGCCCTTGGTCAGGACATGGTGTCAACC
 AGAGGCCACCAACTCTTCTTCTCCTCCTTCTCCTTCCAGCGCGGAAGGCATGTGCGGAGCTAC
 AATCACCTCAAGGAGATGTCCGCTGGAGAAAGCTATTCTCTTCCACCAAGTACTTTCTCAAGATTGAGA
 AGAACGGGAAGGTCAGCGGGACCAAGAAGGAGAAGTCCCGTACAGCATCCTGGAGATAACATCAGTAGA
 AATCGGAGTTGTGCCGTCAAAGCCATTAACAGCAACTATTACTTAGCCATGAACAAGAAGGGGAAACTC
 TATGGCTCAAAAAGAAATTTAAACAATGACTGTAAGCTGAAGGAGAGGATAGAGGAAAATGGATACAATACCT
 ATGCATCATTTAACTGGCAGCATAATGGGAGGCAAATGTATGTGGCATTGAATGGAAAAGGAGCTCCAAG
 GAGAGGACAGAAAACACGAAGGAAAAACACCTCTGCTCACTTTCTCCAATGGTGGTACACTCA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC210120 representing NM_004465
 Red=Cloning site Green=Tags(s)

MWKWILTHCASAFPHLPGCCCCFFLLFLVSSVPVTCQALQDMVSPEATNSSSSSFSSPSSAGRHVRSY
 NHLQGDVWRKLFSTFKYFLKIEKNGKVSCTKENCYPYSILEITSVEIGVVAVKAINSNYYLAMNKKGKL
 YGSKEFNNDCKLKERIEENGYNTYASFNWQHNGRQMYVALNGKGAPRRGQKTRRKNTSAHFLPMVVHS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mg2997_e01.zip



Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_004465

ORF Size: 624 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.


RefSeq: [NM_004465.2](#)

RefSeq Size: 627 bp


RefSeq ORF: 627 bp

Locus ID:	2255
UniProt ID:	O15520
Cytogenetics:	5p12
Domains:	FGF
Protein Families:	Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS, Secreted Protein, Transcription Factors, Transmembrane
Protein Pathways:	MAPK signaling pathway, Melanoma, Pathways in cancer, Regulation of actin cytoskeleton
MW:	23.44 kDa
Gene Summary:	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein exhibits mitogenic activity for keratinizing epidermal cells, but essentially no activity for fibroblasts, which is similar to the biological activity of FGF7. Studies of the mouse homolog of suggested that this gene is required for embryonic epidermal morphogenesis including brain development, lung morphogenesis, and initiation of lim bud formation. This gene is also implicated to be a primary factor in the process of wound healing. [provided by RefSeq, Jul 2008]

Product images:



Circular map for RC210120



Western blot validation of overexpression lysate (Cat# [LY417970]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC210120 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).