

Product datasheet for RC210134L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LTB (NM_002341) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: LTB (NM_002341) Human Tagged ORF Clone Lentiviral Particle

Symbol: LTB

Synonyms: p33; TNFC; TNFSF3; TNLG1C

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_002341

ORF Size: 732 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC210134).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002341.1

RefSeq Size: 894 bp
RefSeq ORF: 735 bp
Locus ID: 4050
UniProt ID: Q06643
Cytogenetics: 6p21.33
Domains: TNF

Protein Families: Druggable Genome, Transmembrane





LTB (NM_002341) Human Tagged ORF Clone Lentiviral Particle - RC210134L4V

Protein Pathways: Cytokine-cytokine receptor interaction

MW: 25.2 kDa

Gene Summary: Lymphotoxin beta is a type II membrane protein of the TNF family. It anchors lymphotoxin-

alpha to the cell surface through heterotrimer formation. The predominant form on the lymphocyte surface is the lymphotoxin-alpha 1/beta 2 complex (e.g. 1 molecule alpha/2 molecules beta) and this complex is the primary ligand for the lymphotoxin-beta receptor. The minor complex is lymphotoxin-alpha 2/beta 1. LTB is an inducer of the inflammatory response system and involved in normal development of lymphoid tissue. Lymphotoxin-beta

isoform b is unable to complex with lymphotoxin-alpha suggesting a function for

lymphotoxin-beta which is independent of lympyhotoxin-alpha. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]