

## Product datasheet for RC210146L4

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Caspase 14 (CASP14) (NM\_012114) Human Tagged Lenti ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: Caspase 14 (CASP14) (NM\_012114) Human Tagged Lenti ORF Clone

Tag: mGFP

Symbol: Caspase 14

Synonyms: ARCI12

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Sgfl-Mlul

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC210146).

Sequence:

Restriction Sites: Cloning Scheme:

Cloning sites used for ORF Shuttling:





st The last codon before the Stop codon of the ORF.

**ACCN:** NM\_012114

ORF Size: 726 bp



#### Caspase 14 (CASP14) (NM\_012114) Human Tagged Lenti ORF Clone - RC210146L4

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 012114.1</u>

 RefSeq Size:
 777 bp

 RefSeq ORF:
 729 bp

 Locus ID:
 23581

 UniProt ID:
 P31944

 Cytogenetics:
 19p13.12

**Protein Families:** Druggable Genome

**MW:** 27.5 kDa

**Gene Summary:** This gene encodes a member of the cysteine-aspartic acid protease (caspase) family.

Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This caspase has been shown to be processed and activated by caspase 8 and caspase 10 in vitro, and by anti-Fas agonist antibody or TNF-related apoptosis inducing ligand


in vivo. The expression and processing of this caspase may be involved in keratinocyte

terminal differentiation, which is important for the formation of the skin barrier. [provided by

RefSeq, Jul 2008]



# **Product images:**



Circular map for RC210146L4