

Product datasheet for RC210193L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CRISP3 (NM_006061) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CRISP3 (NM 006061) Human Tagged ORF Clone Lentiviral Particle

Symbol: CRISP3

Synonyms: Aeg2; CRISP-3; CRS3; dJ442L6.3; SGP28

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_006061

ORF Size: 735 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC210193).

Sequence:
OTI Disclaimer:

Domains:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 006061.1, NP 006052.1

SCP

 RefSeq Size:
 2219 bp

 RefSeq ORF:
 777 bp

 Locus ID:
 10321

 UniProt ID:
 P54108

 Cytogenetics:
 6p12.3

Protein Families: Secreted Protein



ORIGENE

MW: 27.6 kDa

Gene Summary:

This gene encodes a member of the cysteine-rich secretory protein (CRISP) family within the CRISP, antigen 5 and pathogenesis-related 1 proteins superfamily. The encoded protein has an N-terminal CRISP, antigen 5 and pathogenesis-related 1 proteins domain, a hinge region, and a C-terminal ion channel regulator domain. This protein contains cysteine residues, located in both the N- and C-terminal domains, that form eight disulfide bonds, a distinguishing characteristic of this family. This gene is expressed in the male reproductive tract where it plays a role in sperm function and fertilization, and the female reproductive tract where it plays a role in endometrial receptivity for embryo implantation. This gene is upregulated in certain types of prostate cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2016]