

Product datasheet for **RC210197L4V**

Phospholipase C beta 1 (PLCB1) (NM_015192) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Phospholipase C beta 1 (PLCB1) (NM_015192) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Phospholipase C beta 1
Synonyms:	DEE12; EIEE12; PI-PLC; PLC-154; PLC-beta-1; PLC-I; PLC154; PLCB1A; PLCB1B
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_015192
ORF Size:	3648 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC210197).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_015192.2
RefSeq Size:	7103 bp
RefSeq ORF:	3651 bp
Locus ID:	23236
UniProt ID:	Q9NQ66
Cytogenetics:	20p12.3
Domains:	C2, PI-PLC-X, PI-PLC-Y
Protein Families:	Druggable Genome



[View online »](#)

Protein Pathways: Alzheimer's disease, Calcium signaling pathway, Chemokine signaling pathway, Gap junction, GnRH signaling pathway, Huntington's disease, Inositol phosphate metabolism, Long-term depression, Long-term potentiation, Melanogenesis, Metabolic pathways, Phosphatidylinositol signaling system, Vascular smooth muscle contraction, Wnt signaling pathway

MW: 138.6 kDa

Gene Summary: The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of many extracellular signals. This gene is activated by two G-protein alpha subunits, alpha-q and alpha-11. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]