

Product datasheet for RC210230L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EBP1 (PA2G4) (NM_006191) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: EBP1 (PA2G4) (NM 006191) Human Tagged ORF Clone Lentiviral Particle

Symbol: EBP

Synonyms: EBP1; HG4-1; p38-2G4

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_006191 **ORF Size:** 1182 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC210230).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 006191.2</u>

 RefSeq Size:
 2643 bp

 RefSeq ORF:
 1185 bp

 Locus ID:
 5036

 UniProt ID:
 Q9UQ80

 Cytogenetics:
 12q13.2

Domains: Peptidase_M24

Protein Families: Druggable Genome, Protease, Stem cell - Pluripotency





MW: 43.6 kDa

Gene Summary:

This gene encodes an RNA-binding protein that is involved in growth regulation. This protein is present in pre-ribosomal ribonucleoprotein complexes and may be involved in ribosome assembly and the regulation of intermediate and late steps of rRNA processing. This protein can interact with the cytoplasmic domain of the ErbB3 receptor and may contribute to transducing growth regulatory signals. This protein is also a transcriptional co-repressor of androgen receptor-regulated genes and other cell cycle regulatory genes through its interactions with histone deacetylases. This protein has been implicated in growth inhibition and the induction of differentiation of human cancer cells. Six pseudogenes, located on chromosomes 3, 6, 9, 18, 20 and X, have been identified. [provided by RefSeq, Jul 2008]