

## Product datasheet for RC210231L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CYP2C18 (NM\_000772) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: CYP2C18 (NM 000772) Human Tagged ORF Clone Lentiviral Particle

Symbol: CYP2C18

**Synonyms:** CPCI; CYP2C; CYP2C17; P450-6B/29C; P450IIC17

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_000772

 ORF Size:
 1470 bp

OTT 5126.

ORF Nucleotide Sequence:

The ORF insert of this clone is exactly the same as(RC210231).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000772.1

 RefSeq Size:
 1995 bp

 RefSeq ORF:
 1473 bp

 Locus ID:
 1562

 UniProt ID:
 P33260

 Cytogenetics:
 10q23.33

**Domains:** p450

**Protein Families:** Druggable Genome, P450





## CYP2C18 (NM\_000772) Human Tagged ORF Clone Lentiviral Particle - RC210231L3V

**Protein Pathways:** Arachidonic acid metabolism, Drug metabolism - cytochrome P450, Linoleic acid metabolism,

Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Retinol metabolism

**MW:** 55.5 kDa

**Gene Summary:** This gene encodes a member of the cytochrome P450 superfamily of enzymes. The

cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum but its specific substrate has not yet been determined. The gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. An additional gene, CYP2C17, was once thought to exist; however, CYP2C17 is now considered an artefact based on a chimera of CYP2C18 and CYP2C19. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul

2008]