

Product datasheet for RC210356L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PON1 (NM_000446) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PON1 (NM_000446) Human Tagged ORF Clone Lentiviral Particle

Symbol: PON'

Synonyms: ESA; MVCD5; PON

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_000446 **ORF Size:** 1065 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC210356).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000446.3

 RefSeq Size:
 1769 bp

 RefSeq ORF:
 1068 bp

 Locus ID:
 5444

 UniProt ID:
 P27169

 Cytogenetics:
 7q21.3

Domains: Arylesterase

Protein Families: Druggable Genome, Secreted Protein





PON1 (NM_000446) Human Tagged ORF Clone Lentiviral Particle - RC210356L2V

Protein Pathways: Metabolic pathways

MW: 39.7 kDa

Gene Summary: This gene encodes a member of the paraoxonase family of enzymes and exhibits lactonase

and ester hydrolase activity. Following synthesis in the kidney and liver, the enzyme is secreted into the circulation, where it binds to high density lipoprotein (HDL) particles and hydrolyzes thiolactones and xenobiotics, including paraoxon, a metabolite of the insecticide parathion. Polymorphisms in this gene may be associated with coronary artery disease and diabetic retinopathy. The gene is found in a cluster of three related paraoxonase genes on

chromosome 7. [provided by RefSeq, Aug 2017]