

## Product datasheet for RC210361L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Galactoside 2 alpha L fucosyltransferase 1 (FUT1) (NM\_000148) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Galactoside 2 alpha L fucosyltransferase 1 (FUT1) (NM\_000148) Human Tagged ORF Clone

Lentiviral Particle

Symbol: Galactoside 2 alpha L fucosyltransferase 1

Synonyms: H; HH; HSC

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM\_000148

**ORF Size:** 1095 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC210361).

**OTI Disclaimer:** 

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 000148.2</u>

 RefSeq Size:
 4246 bp

 RefSeq ORF:
 1098 bp

 Locus ID:
 2523

 UniProt ID:
 P19526

 Cytogenetics:
 19q13.33

**Protein Families:** Druggable Genome, Transmembrane





## Galactoside 2 alpha L fucosyltransferase 1 (FUT1) (NM\_000148) Human Tagged ORF Clone Lentiviral Particle – RC210361L3V

**Protein Pathways:** Glycosphingolipid biosynthesis - globo series, Glycosphingolipid biosynthesis - lacto and

neolacto series, Metabolic pathways

MW: 41.3 kDa

**Gene Summary:** This gene encodes a Golgi stack membrane protein that is involved in the creation of a

precursor of the H antigen, which is required for the final step in the synthesis of soluble A and B antigens. This is one of two genes encoding the galactoside 2-L-fucosyltransferase enzyme. Mutations in this gene are a cause of the H-Bombay blood group. [provided by

RefSeq, Aug 2016]