

Product datasheet for RC210371L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MS4A2 (NM_000139) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: MS4A2 (NM 000139) Human Tagged ORF Clone Lentiviral Particle

Symbol: MS4A2

Synonyms: APY; ATOPY; FCER1B; FCERI; IGEL; IGER; IGHER; MS4A1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 000139

ORF Size: 732 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC210371).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000139.2

 RefSeq Size:
 3658 bp

 RefSeq ORF:
 735 bp

 Locus ID:
 2206

 UniProt ID:
 Q01362

 Cytogenetics:
 11q12.1

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Asthma, Fc epsilon RI signaling pathway





ORIGENE

MW: 26.5 kDa

Gene Summary:

The allergic response involves the binding of allergen to receptor-bound IgE followed by cell activation and the release of mediators responsible for the manifestations of allergy. The IgE-receptor, a tetramer composed of an alpha, beta, and 2 disulfide-linked gamma chains, is found on the surface of mast cells and basophils. This gene encodes the beta subunit of the high affinity IgE receptor which is a member of the membrane-spanning 4A gene family. Members of this nascent protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. This family member is localized to 11q12, among a cluster of membrane-spanning 4A gene family members. Alternative splicing results in multiple transcript variants encoding distinct proteins. Additional transcript variants have been described but require experimental validation. [provided by RefSeq, Mar 2012]