

## Product datasheet for RC210395L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Neurokinin 1 Receptor (TACR1) (NM\_001058) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: Neurokinin 1 Receptor (TACR1) (NM\_001058) Human Tagged ORF Clone Lentiviral Particle

Symbol: Neurokinin 1 Receptor
Synonyms: NK1R; NKIR; SPR; TAC1R

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001058 **ORF Size:** 1221 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC210395).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001058.2</u>

 RefSeq Size:
 4795 bp

 RefSeq ORF:
 1224 bp

 Locus ID:
 6869

 UniProt ID:
 P25103

 Cytogenetics:
 2p12

 Domains:
 7tm 1

**Protein Families:** Druggable Genome, GPCR, Transmembrane





## Neurokinin 1 Receptor (TACR1) (NM\_001058) Human Tagged ORF Clone Lentiviral Particle – RC210395L4V

**Protein Pathways:** Calcium signaling pathway, Neuroactive ligand-receptor interaction

MW: 46.3 kDa

**Gene Summary:** This gene belongs to a gene family of tachykinin receptors. These tachykinin receptors are

characterized by interactions with G proteins and contain seven hydrophobic

transmembrane regions. This gene encodes the receptor for the tachykinin substance P, also

referred to as neurokinin 1. The encoded protein is also involved in the mediation of phosphatidylinositol metabolism of substance P. [provided by RefSeq, Sep 2008]