

## Product datasheet for RC210406L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## P2Y1 (P2RY1) (NM 002563) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type: Lentiviral Particles** 

**Product Name:** P2Y1 (P2RY1) (NM\_002563) Human Tagged ORF Clone Lentiviral Particle

Symbol:

P2Y1; SARCC Synonyms:

**Mammalian Cell** 

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

mGFP Tag:

NM 002563 ACCN: **ORF Size:** 1119 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC210406).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 002563.2

RefSeq Size: 3110 bp RefSeq ORF: 1122 bp Locus ID: 5028 **UniProt ID:** P47900 Cytogenetics: 3q25.2 **Domains:** 

**Protein Families:** Druggable Genome, GPCR, Transmembrane

7tm 1





## P2Y1 (P2RY1) (NM\_002563) Human Tagged ORF Clone Lentiviral Particle - RC210406L2V

**Protein Pathways:** Neuroactive ligand-receptor interaction

MW: 42.1 kDa

**Gene Summary:** The product of this gene belongs to the family of G-protein coupled receptors. This family has

several receptor subtypes with different pharmacological selectivity, which overlaps in some cases, for various adenosine and uridine nucleotides. This receptor functions as a receptor

for extracellular ATP and ADP. In platelets binding to ADP leads to mobilization of

intracellular calcium ions via activation of phospholipase C, a change in platelet shape, and

probably to platelet aggregation. [provided by RefSeq, Jul 2008]