

### Product datasheet for RC210452L4V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Proteasome subunit beta type 2 (PSMB2) (NM\_002794) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** Proteasome subunit beta type 2 (PSMB2) (NM\_002794) Human Tagged ORF Clone Lentiviral

Particle

**Symbol:** Proteasome subunit beta type 2

Synonyms: HC7-I

Mammalian Cell Pure

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_002794

ORF Size: 603 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC210452).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 002794.3</u>

 RefSeq Size:
 4759 bp

 RefSeq ORF:
 606 bp

 Locus ID:
 5690

 UniProt ID:
 P49721

 Cytogenetics:
 1p34.3

**Domains:** proteasome





## Proteasome subunit beta type 2 (PSMB2) (NM\_002794) Human Tagged ORF Clone Lentiviral Particle - RC210452L4V

**Protein Families:** Druggable Genome, Protease

Protein Pathways: Proteasome MW: 22.8 kDa

**Gene Summary:** The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S

core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. Multiple alternatively spliced transcript variants encoding

distinct isoforms have been found for this gene. [provided by RefSeq, Dec 2010]